

AD-A148 664

THE ADSORPTION OF ACRYLONITRILE AT A GOLD ELECTRODE: AN
EMIRS MEASUREMENT. (U) UTAH UNIV SALT LAKE CITY DEPT OF
CHEMISTRY A BEWICK ET AL. 28 NOV 84 TR-39

1/1

UNCLASSIFIED

N00014-83-K-0470


F/G 7/4

NL

END

FILMED

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A

(12)

OFFICE OF NAVAL RESEARCH

Contract N00014-83-K-0470

Task No. NR 359-718

TECHNICAL REPORT NO. 39

AD-A148 664

The Adsorption of Acrylonitrile at a Gold Electrode:
An EMIRS Measurement of the Electrochemical Stark Effect

By

A. Bewick
C. Gibilaro
Stanley Pons

Prepared for Publication in
Langmuir

University of Utah
Department of Chemistry
Salt Lake City, Utah 84112

DTIC
ELECTE
DEC 21 1984
S B

November 28, 1984

Reproduction in whole or in part is permitted for
any purpose of the United States Government.

This document has been approved for public release
and sale; its distribution is unlimited.

DTIC FILE COPY

84 12 12 048

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 39	2. GOVT ACCESSION NO. 7D-A148 664	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) The Adsorption of Acrylonitrile at a Gold Electrode: An EMIRS Measurement of the Electrochemical Stark Effect		5. TYPE OF REPORT & PERIOD COVERED Technical Report # 39
7. AUTHOR(s) A. Bewick; C. Gibilaro; S. Pons		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Utah Department of Chemistry Salt Lake City, UT 84112		8. CONTRACT OR GRANT NUMBER(s) N00014-83-K-0470
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Chemistry Program - Chemistry Code 472 Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Task No. NR 359-718
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE November 28, 1984
		13. NUMBER OF PAGES 15
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Acrylonitrile, Infrared Spectroelectrochemistry p.c.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Electrochemically modulated infra red spectroscopy of acrylonitrile adsorbed on a gold electrode in aqueous sulphuric acid shows potential - dependent unipolar absorption bands at $2130/\text{cm}^{-1}$ and $1520/\text{cm}^{-1}$. These are ascribed to heavily red shifted $\text{C}=\text{N}^-$ and $\text{C}=\text{C}$ stretches from acrylonitrile adsorbed flat on the surface by means of the π -electron system. These bands are forbidden by the surface selection rule but are made visible by an Electrochemical Stark Effect. 1		

1 The adsorption of acrylonitrile at a gold electrode: an EMIRS measurement
of the Electrochemical Stark Effect

A. Bewick* and C. Gibilaro
Department of Chemistry,
University of Southampton,
Southampton, SO9 5NH,
England.

and

S. Pons,
Department of Chemistry,
University of Utah,
Salt Lake City,
Utah 84112,
U.S.A.

*To whom correspondence should be addressed.

Abstract

Electrochemically modulated infra red spectroscopy of acrylonitrile adsorbed on a gold electrode in aqueous sulphuric acid shows potential - dependent unipolar absorption bands at 2130 cm^{-1} and 1520 cm^{-1} . These are ascribed to heavily red shifted C N^+ and $\text{C}=\text{C}$ stretches from acrylonitrile adsorbed flat on the surface by means of the π -electron system. These bands are forbidden by the surface selection rule but are made visible by an Electrochemical Stark Effect.



Accession For	
ATIS 10001	<input checked="" type="checkbox"/>
10001	<input type="checkbox"/>
10001	<input type="checkbox"/>
J. 10001	
B. 10001	
D. 10001	
A. 10001	
C. 10001	
E. 10001	
F. 10001	
G. 10001	
H. 10001	
I. 10001	
J. 10001	
K. 10001	
L. 10001	
M. 10001	
N. 10001	
O. 10001	
P. 10001	
Q. 10001	
R. 10001	
S. 10001	
T. 10001	
U. 10001	
V. 10001	
W. 10001	
X. 10001	
Y. 10001	
Z. 10001	
AA. 10001	
AB. 10001	
AC. 10001	
AD. 10001	
AE. 10001	
AF. 10001	
AG. 10001	
AH. 10001	
AI. 10001	
AJ. 10001	
AK. 10001	
AL. 10001	
AM. 10001	
AN. 10001	
AO. 10001	
AP. 10001	
AQ. 10001	
AR. 10001	
AS. 10001	
AT. 10001	
AU. 10001	
AV. 10001	
AW. 10001	
AX. 10001	
AY. 10001	
AZ. 10001	
BA. 10001	
BB. 10001	
BC. 10001	
BD. 10001	
BE. 10001	
BF. 10001	
BG. 10001	
BH. 10001	
BI. 10001	
BJ. 10001	
BK. 10001	
BL. 10001	
BM. 10001	
BN. 10001	
BO. 10001	
BP. 10001	
BQ. 10001	
BR. 10001	
BS. 10001	
BT. 10001	
BU. 10001	
BV. 10001	
BW. 10001	
BX. 10001	
BY. 10001	
BZ. 10001	
CA. 10001	
CB. 10001	
CC. 10001	
CD. 10001	
CE. 10001	
CF. 10001	
CG. 10001	
CH. 10001	
CI. 10001	
CJ. 10001	
CK. 10001	
CL. 10001	
CM. 10001	
CN. 10001	
CO. 10001	
CP. 10001	
CQ. 10001	
CR. 10001	
CS. 10001	
CT. 10001	
CU. 10001	
CV. 10001	
CW. 10001	
CX. 10001	
CY. 10001	
CZ. 10001	
DA. 10001	
DB. 10001	
DC. 10001	
DD. 10001	
DE. 10001	
DF. 10001	
DG. 10001	
DH. 10001	
DI. 10001	
DJ. 10001	
DK. 10001	
DL. 10001	
DM. 10001	
DN. 10001	
DO. 10001	
DP. 10001	
DQ. 10001	
DR. 10001	
DS. 10001	
DT. 10001	
DU. 10001	
DV. 10001	
DW. 10001	
DX. 10001	
DY. 10001	
DZ. 10001	
EA. 10001	
EB. 10001	
EC. 10001	
ED. 10001	
EE. 10001	
EF. 10001	
EG. 10001	
EH. 10001	
EI. 10001	
EJ. 10001	
EK. 10001	
EL. 10001	
EM. 10001	
EN. 10001	
EO. 10001	
EP. 10001	
EQ. 10001	
ER. 10001	
ES. 10001	
ET. 10001	
EU. 10001	
EV. 10001	
EW. 10001	
EX. 10001	
EY. 10001	
EZ. 10001	
FA. 10001	
FB. 10001	
FC. 10001	
FD. 10001	
FE. 10001	
FF. 10001	
FG. 10001	
FH. 10001	
FI. 10001	
FJ. 10001	
FK. 10001	
FL. 10001	
FM. 10001	
FN. 10001	
FO. 10001	
FP. 10001	
FQ. 10001	
FR. 10001	
FS. 10001	
FT. 10001	
FU. 10001	
FV. 10001	
FW. 10001	
FX. 10001	
FY. 10001	
FZ. 10001	
GA. 10001	
GB. 10001	
GC. 10001	
GD. 10001	
GE. 10001	
GF. 10001	
GG. 10001	
GH. 10001	
GI. 10001	
GJ. 10001	
GK. 10001	
GL. 10001	
GM. 10001	
GN. 10001	
GO. 10001	
GP. 10001	
GQ. 10001	
GR. 10001	
GS. 10001	
GT. 10001	
GU. 10001	
GV. 10001	
GW. 10001	
GX. 10001	
GY. 10001	
GZ. 10001	
HA. 10001	
HB. 10001	
HC. 10001	
HD. 10001	
HE. 10001	
HF. 10001	
HG. 10001	
HH. 10001	
HI. 10001	
HJ. 10001	
HK. 10001	
HL. 10001	
HM. 10001	
HN. 10001	
HO. 10001	
HP. 10001	
HQ. 10001	
HR. 10001	
HS. 10001	
HT. 10001	
HU. 10001	
HV. 10001	
HW. 10001	
HX. 10001	
HY. 10001	
HZ. 10001	
IA. 10001	
IB. 10001	
IC. 10001	
ID. 10001	
IE. 10001	
IF. 10001	
IG. 10001	
IH. 10001	
II. 10001	
IJ. 10001	
IK. 10001	
IL. 10001	
IM. 10001	
IN. 10001	
IO. 10001	
IP. 10001	
IQ. 10001	
IR. 10001	
IS. 10001	
IT. 10001	
IU. 10001	
IV. 10001	
IW. 10001	
IX. 10001	
IY. 10001	
IZ. 10001	
JA. 10001	
JB. 10001	
JC. 10001	
JD. 10001	
JE. 10001	
JF. 10001	
JG. 10001	
JH. 10001	
JI. 10001	
JJ. 10001	
JK. 10001	
JL. 10001	
JM. 10001	
JN. 10001	
JO. 10001	
JP. 10001	
JQ. 10001	
JR. 10001	
JS. 10001	
JT. 10001	
JU. 10001	
JV. 10001	
JW. 10001	
JX. 10001	
JY. 10001	
JZ. 10001	
KA. 10001	
KB. 10001	
KC. 10001	
KD. 10001	
KE. 10001	
KF. 10001	
KG. 10001	
KH. 10001	
KI. 10001	
KJ. 10001	
KK. 10001	
KL. 10001	
KM. 10001	
KN. 10001	
KO. 10001	
KP. 10001	
KQ. 10001	
KR. 10001	
KS. 10001	
KT. 10001	
KU. 10001	
KV. 10001	
KW. 10001	
KX. 10001	
KY. 10001	
KZ. 10001	
LA. 10001	
LB. 10001	
LC. 10001	
LD. 10001	
LE. 10001	
LF. 10001	
LG. 10001	
LH. 10001	
LI. 10001	
LJ. 10001	
LK. 10001	
LL. 10001	
LM. 10001	
LN. 10001	
LO. 10001	
LP. 10001	
LQ. 10001	
LR. 10001	
LS. 10001	
LT. 10001	
LU. 10001	
LV. 10001	
LW. 10001	
LX. 10001	
LY. 10001	
LZ. 10001	
MA. 10001	
MB. 10001	
MC. 10001	
MD. 10001	
ME. 10001	
MF. 10001	
MG. 10001	
MH. 10001	
MI. 10001	
MJ. 10001	
MK. 10001	
ML. 10001	
MM. 10001	
MN. 10001	
MO. 10001	
MP. 10001	
MQ. 10001	
MR. 10001	
MS. 10001	
MT. 10001	
MU. 10001	
MV. 10001	
MW. 10001	
MX. 10001	
MY. 10001	
MZ. 10001	
NA. 10001	
NB. 10001	
NC. 10001	
ND. 10001	
NE. 10001	
NF. 10001	
NG. 10001	
NH. 10001	
NI. 10001	
NJ. 10001	
NK. 10001	
NL. 10001	
NM. 10001	
NN. 10001	
NO. 10001	
NP. 10001	
NQ. 10001	
NR. 10001	
NS. 10001	
NT. 10001	
NU. 10001	
NV. 10001	
NW. 10001	
NX. 10001	
NY. 10001	
NZ. 10001	
OA. 10001	
OB. 10001	
OC. 10001	
OD. 10001	
OE. 10001	
OF. 10001	
OG. 10001	
OH. 10001	
OI. 10001	
OJ. 10001	
OK. 10001	
OL. 10001	
OM. 10001	
ON. 10001	
OO. 10001	
OP. 10001	
OQ. 10001	
OR. 10001	
OS. 10001	
OT. 10001	
OU. 10001	
OV. 10001	
OW. 10001	
OX. 10001	
OY. 10001	
OZ. 10001	
PA. 10001	
PB. 10001	
PC. 10001	
PD. 10001	
PE. 10001	
PF. 10001	
PG. 10001	
PH. 10001	
PI. 10001	
PJ. 10001	
PK. 10001	
PL. 10001	
PM. 10001	
PN. 10001	
PO. 10001	
PP. 10001	
PQ. 10001	
PR. 10001	
PS. 10001	
PT. 10001	
PU. 10001	
PV. 10001	
PW. 10001	
PX. 10001	
PY. 10001	
PZ. 10001	
QA. 10001	
QB. 10001	
QC. 10001	
QD. 10001	
QE. 10001	
QF. 10001	
QG. 10001	
QH. 10001	
QI. 10001	
QJ. 10001	
QK. 10001	
QL. 10001	
QM. 10001	
QN. 10001	
QO. 10001	
QP. 10001	
QQ. 10001	
QR. 10001	
QS. 10001	
QT. 10001	
QU. 10001	
QV. 10001	
QW. 10001	
QX. 10001	
QY. 10001	
QZ. 10001	
RA. 10001	
RB. 10001	
RC. 10001	
RD. 10001	
RE. 10001	
RF. 10001	
RG. 10001	
RH. 10001	
RI. 10001	
RJ. 10001	
RK. 10001	
RL. 10001	
RM. 10001	
RN. 10001	
RO. 10001	
RP. 10001	
RQ. 10001	
RR. 10001	
RS. 10001	
RT. 10001	
RU. 10001	
RV. 10001	
RW. 10001	
RX. 10001	
RY. 10001	
RZ. 10001	
SA. 10001	
SB. 10001	
SC. 10001	
SD. 10001	
SE. 10001	
SF. 10001	
SG. 10001	
SH. 10001	
SI. 10001	
SJ. 10001	
SK. 10001	
SL. 10001	
SM. 10001	
SN. 10001	
SO. 10001	
SP. 10001	
SQ. 10001	
SR. 10001	
SS. 10001	
ST. 10001	
SU. 10001	
SV. 10001	
SW. 10001	
SX. 10001	
SY. 10001	
SZ. 10001	
TA. 10001	
TB. 10001	
TC. 10001	
TD. 10001	
TE. 10001	
TF. 10001	
TG. 10001	
TH. 10001	
TI. 10001	
TJ. 10001	
TK. 10001	
TL. 10001	
TM. 10001	
TN. 10001	
TO. 10001	
TP. 10001	
TQ. 10001	
TR. 10001	
TS. 10001	
TT. 10001	
TU. 10001	
TV. 10001	
TW. 10001	
TX. 10001	
TY. 10001	
TZ. 10001	
UA. 10001	
UB. 10001	
UC. 10001	
UD. 10001	
UE. 10001	
UF. 10001	
UG. 10001	
UH. 10001	
UI. 10001	
UJ. 10001	
UK. 10001	
UL. 10001	
UM. 10001	
UN. 10001	
UO. 10001	
UP. 10001	
UQ. 10001	
UR. 10001	
US. 10001	
UT. 10001	
UU. 10001	
UV. 10001	
UW. 10001	
UX. 10001	
UY. 10001	
UZ. 10001	
VA. 10001	
VB. 10001	
VC. 10001	
VD. 10001	
VE. 10001	
VF. 10001	
VG. 10001	
VH. 10001	
VI. 10001	
VJ. 10001	
VK. 10001	
VL. 10001	
VM. 10001	
VN. 10001	
VO. 10001	
VP. 10001	
VQ. 10001	
VR. 10001	
VS. 10001	
VT. 10001	
VU. 10001	
VV.	

Introduction

The newly developed methods for obtaining in situ the infra red spectra from species adsorbed on electrode surfaces are finding a wide range of applications [1, 2, 3, 4, 5]. A particularly useful feature of these methods is their sensitivity towards the orientation of an adsorbed molecule with respect to the metal surface. The surface selection rule predicts that only those vibrational modes of the molecule which possess a non-zero component of the dipole derivative in a direction normal to the surface are able to interact with the infra red radiation reflected from the surface, and the intensities of the observed bands will be proportional to the cosine of the angle between the surface and the normal mode coordinate dipole derivative from that band. This requirement has been used to infer the orientation of benzonitrile on a gold electrode [5, 6] and difluorobenzene [7, 8] on a platinum electrode. The former was shown to bond via the nitrogen in a configuration normal to the surface, whereas the latter was adsorbed with the aromatic ring flat on the surface except at high concentrations when a skewed configuration was adopted.

The acrylonitrile molecule is in some ways similar to benzonitrile and it possesses orbitals which could allow interaction with a metal surface either lying flat on the surface or sitting up with the nitrogen bonded to the metal. An investigation was carried out using electrochemically modulated infra red spectroscopy (EMIRS) to deduce its orientation adsorbed on a gold electrode, this being part of a series of investigations to determine the electronic factors, both in the adsorbate and the adsorbant, which are important in determining their relative orientations.

Experimental

Solutions were prepared using water distilled four times: a first distillation, without additives, from glass then slowly a second time after addition of a little alkaline potassium permanganate; the third distillation followed after addition of a little orthophosphoric acid and a final distillation without additives was made immediately prior to the experiment. Aristar grade sulphuric acid was used without further purification to prepare the IM solutions used in all experiments. Acrylonitrile (BDH) was freshly distilled before use to eliminate the trace of p-methoxyphenol added to inhibit polymerisation. All glassware was cleaned by prolonged immersion in a mixture of equal volumes of concentrated nitric and sulphuric acids followed by rinsing with distilled water and a final boiling for 30 minutes with triply distilled water. All solutions were deaerated in the spectroelectrochemical cell with a nitrogen purge for 30 minutes.

The gold electrode was polished to a mirror finish using alumina of decreasing grades down to $0.05\mu\text{m}$ on a water lubricated polishing cloth. The cell was fitted with a single crystal, intrinsic silicon window and the angle of incidence (to the normal) was set at just below 70° . The spectrometer and associated equipment were as already describe [1]. The modulation frequency of the electrode potential was set to 8.5 Hz and a Pye IR 50 detector was employed, this detector being about twenty times less sensitive than the mercury cadmium telluride detector currently used. The values of relative reflectivity change, $\Delta R/R$, plotted are equivalent to absorbance units and the sign convention adopted produces positive bands for increased absorption of the radiation at the more negative (or less positive) potential. All electrode potentials are quoted against the saturated calomel electrode. The control and modulation of the electrode

potential was provided by a Hi-Tek Instruments Potentiostat Type DT2101 and Waveform Generator Type PPR1.

Results and Discussion

Linear sweep voltammograms of the gold electrode in 1M H_2SO_4 with and without added acrylonitrile in the range 0.05 to 0.5M indicated adsorption of the organic. This was confirmed by differential capacity measurements carried out at various a.c. modulation frequencies and various concentrations of acrylonitrile. An example using a frequency of 8.5 Hz is shown in figure 1. This shows a considerable lowering of the differential capacity over the whole of the potential range employed which indicates strong adsorption of the acrylonitrile; the sharp rise beyond IV is evidence for desorption and the onset of oxide formation on the gold. At high concentrations of acrylonitrile (0.4 M) time dependent phenomena were observed which were probably caused by polymerization on the surface [9]. This was not observed at the much lower concentrations used in all the EMIRS measurements nor were any infra red bands seen at 2115 cm^{-1} or 2165 cm^{-1} corresponding to polymer formation [9]. The capacitance plots indicate that bisulphate ion is probably also adsorbed; note the humps at 0.580V and 0.850V. It is apparent also that it is not possible to define modulation potential limits to produce a clear cut adsorption/desorption process free from the effects of oxide formation or hydrogen evolution.

EMIRS measurements for a range of modulation potentials covering the range from -0.2V to +0.95V were carried out over the wavenumber interval from 1950 cm^{-1} to 2450 cm^{-1} . Two bands were observed at about 2130 cm^{-1} and 1520 cm^{-1} the amplitudes and positions of which were potential dependent. Examples of these are shown in figures 2 and 3. The large noise amplitude on the lower wavenumber spectra are due to the low energy throughput in the region of the water bending mode absorption band at 1600

cm^{-1} following strong absorption in the thin aqueous layer between the electrode and the cell window. These two bands are the only features observed and it is interesting to note that they would correspond to the $\text{C}\equiv\text{N}$ and $\text{C}=\text{C}$ stretching modes of adsorbed acrylonitrile if the bonding to the surface lowered each of these frequencies by about 100 cm^{-1} compared to the vibrations of the free molecule. If the molecule was adsorbed via the nitrogen, the frequency shifts would be expected to be much smaller (particularly for the $\text{C}=\text{C}$ stretch) and the $\text{C}\equiv\text{N}$ stretch should shift in the opposite direction. Thus the band positions indicate flat adsorption of the molecule with bonding to the surface via the conjugated π -system. At first sight, however, this conjecture appears to lead to a number of logical difficulties:

- (i) the observed bands are single sided whereas bipolar bands are to be expected (see for example adsorbed CO on Pt [10, 11]) from an adsorbed layer possessing approximately the same surface coverage at each potential defined by the modulation and made observable because the band positions are potential dependent; if the coverage does change with potential, then additional bands of opposite sign and at the frequencies of the solution-free molecule should be, present since adsorption leads to depletion of the solution, but these are not seen; moreover, the largest amplitude for all of the bands should be observed for modulation extending to the most positive potential where desorption should be greatest, however, the opposite effect is seen (see later);
- (ii) the surface selection rule predicts that the $\text{C}\equiv\text{N}$ and $\text{C}=\text{C}$ stretches for flatly adsorbed acrylonitrile should not be observable since, in each case, the dipole derivative with respect to the normal mode coordinate should be entirely parallel to the metal surface

(neglecting the effects of slight changes in symmetry brought about by the adsorbate/adsorbent bonding);

- (iii) the bands are weak, the amplitude of the 2130 cm^{-1} band is an order of magnitude smaller than the C N stretch band observed for a monolayer of adsorbed benzonitrile at a gold electrode [5, 6];
- (iv) the 1520 cm^{-1} band is stronger than the 2130 cm^{-1} band, whereas the C=C and C \equiv N stretches for the free molecule shows the opposite relative amplitudes.

It appears, therefore, that the acrylonitrile molecule is adsorbed flat on the gold surface but that a special mechanism is operating to produce relatively small amplitude unipolar bands i.e. that the two normal mode vibrations produce a dipole derivative vector at an angle to the surface. There are two possible mechanisms which could lead to this effect:

- (a) the symmetry and bonding in the adsorbate/adsorbent complex produces nuclear motions leading to a finite component of the dipole derivative normal to the surface;
- (b) there is vibronic coupling, via the molecular polarisability tensor, to the very high static electrical field in the double layer at the adsorption sites [3, 5, 12, 13, 14].

Using the broadest definition of the Stark Effect, mechanism (b) is an Electrochemical Stark Effect. It is also related to the coupling observed between vibrational modes in charge transfer complexes and the extent of charge transfer; this has been shown to lead to greatly enhanced infra red absorbances in a number of systems [15, 16]. The potential dependence of the amplitude of the 2130 cm^{-1} band, figure 4, shows clearly that the Stark Effect mechanism is the more likely of the two alternatives and thus the present example is the first recognised occurrence of the

Electrochemical Stark Effect. A simple explanation of the effect is that the field in the double layer will induce a dipole in the molecule in a direction normal to the electrode surface the magnitude of which is proportional to the molecular polarisability in that direction. During a vibration the polarisability will vary with the internuclear separations and thus the induced dipole will have a fluctuating component which allows the vibrational mode to interact with radiation. In a first approximation, the Electrochemical Stark Effect requires that the integrated intensity of the infra red absorption brought about by the effect should be proportional to the square of the strength of the static electrical field [12, 17]. It is difficult to test this exactly in the present case because the potential at which the field strength is zero is not known nor is the potential dependence of the field. However, for modulation straddling the zero point there should be a maximum in the measured intensity when one of the modulation limits is at the zero field potential. Figure 4 does in fact show a maximum at a potential of about +0.7V; this would not be an unreasonable position for the potential of zero field. The shape of figure 4 cannot be reconciled with mechanism (a) nor, as mentioned earlier, can it be caused by a potential dependent adsorption/desorption process. The dotted line sketched on figure 4 shows the shape of the potential dependence expected for a square law relationship. In this calculation it has been assumed that the field strength in the inner region of the double layer is proportional to the electrode potential and with the zero located at +0.7V. This direct proportionality can only be a very rough approximation in view of the large variation in the total double layer capacity shown in figure 1 and also the likely occurrence of HSO_4^- adsorption. Bearing these restrictions in mind, the fit is sufficiently good to support our

explanation. The potential dependence of the band position, is also interesting, figure 5. Unlike the data for other simpler examples, such as adsorbed CO on Pt or CN^- on Ag [11, 18], it is non linear. In the present case, if adsorption involves some transfer of electrons to the metal due to overlap of the occupied π -orbitals with metallic orbitals, the shift with potential would be expected to be in the opposite direction to that observed. Thus the observed potential dependence is probably a further consequence of the Stark Effect although it is not calculable without making unwarrantable assumptions.

We conclude, therefore, that the example we have discussed shows the Electrochemical Stark Effect and examination of some of our earlier data for other systems also shows evidence for the operation of this mechanism. The effect has advantages and disadvantages for these studies of adsorption: on the one hand, it provides a tool for probing electrical fields in the double layer and for seeing species otherwise invisible due to the surface selection rule; on the other hand, it makes all the Raman modes of a molecule also IR active and therefore places some additional uncertainties on deductions of adsorbate orientation.

Acknowledgements

Support for this work from the Office of Naval Research Washington, the SERC and a NATO Research Grant (204.83) is gratefully acknowledged.


References


1. A. Bewick, K. Kunimatsu, B.S. Pons and J.W. Russell, J. Electroanal. Chem., 160 (1984) 47.
2. S. Pons, T. Davidson and A. Bewick, J. Electroanal. Chem., 160 (1984)
3. A. Bewick, J. Electroanal. Chem., 150 (1983) 481.
4. S. Pons, J. Electroanal. Chem., 150 (1983) 495.
5. A. Bewick and S.B. Pons, Chapter in "Advances in Infrared and Raman Spectroscopy" eds. R. Hester and R.E. Clarke, 1984, in press.
6. A. Bewick, J. Hudec and M. Razaq, to be submitted.
7. S. Pons and A. Bewick, Langmuir, submitted.
8. A. Bewick and S.B. Pons, Chapter in "Electroanalytical Chemistry", ed. A.J. Bard, submitted.
9. A.Z. Trifinov and I.D. Schopov, J. Electroanal. Chem., 35 (1972) 415.
10. B. Beden, A. Bewick, K. Kunimatsu and C. Lamy, J. Electroanal. Chem., 142 (1982) 345.
11. J.W. Russell, M. Severson, K. Scanlon, J. Overend and A. Bewick, J. Phys. Chem., 87 (1983) 293.
12. S. Pons, A. Bewick, and C. Korzeniewski, J. Phys. Chem., submitted.
13. A. Bewick, S. Pons and C. Korzeniewski, Electrochim. Acta, submitted.
14. S. Pons, S.B. Khoo, A. Bewick, M. Datta, J.J. Smith, A.S. Hinman and G. Zachmann, J. Phys. Chem., 88 (1984) 3575.

15. J. P. Devlin, and K. Consani, J. Phys. Chem. 85(1981)2597.
16. E.E. Ferguson and F.A. Matson, J. Amer. Chem. Soc., 82(1960)3268.
17. M.F. Crawford and R.E. MacDonald, Can. J. Phys. 36(1958)102
18. A. Bewick and C. Gibilaro, in preparation.


Legends for Figures

- Figure 1. Differential capacity versus potential plots for a gold electrode in (a) 1M H_2SO_4 and (b) 1M H_2SO_4 + 0.05M acrylonitrile; a.c. modulation 5 mV peak to peak at 8.5 Hz superimposed on a linear sweep at 10^{-2} V s^{-1} .
- Figure 2. EMIRS spectra from a gold electrode in 1M H_2SO_4 + 0.05M acrylonitrile. Modulation at 8.5 Hz from -0.195 V to (a) +0.750 V, (b) +0.350 V and (c) +0.050 V.
- Figure 3. EMIRS spectrum from a gold electrode in 1M H_2SO_4 + 0.05M acrylonitrile. Modulation at 8.5 Hz from -0.195 V to +0.750 V.
- Figure 4. Potential dependence of the intensity of the $\text{C}\equiv\text{N}$ stretch band from adsorbed acrylonitrile. The dashed curve illustrates a calculated square law field dependence.
- Figure 5. Potential dependence of the position of the $\text{C}\equiv\text{N}$ stretch band from adsorbed acrylonitrile.







10⁻² 25.7



cm⁻¹



114.5



END

FILMED

1-85

DTIC